

20% BABY PIG TI35C400

MEDICATED

Complete Swine Starter to Increase the Digestibility of Phytin-Bound Phosphorus in Swine Diets

Control of swine dysentery associated with *Brachyspira hyodysenteriae* susceptible to Tiamulin. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENTS

CHLORTETRACYCLINE	400
g/ton TIAMULIN HYDROGEN FUMERATE	
.....	35 g/ton
CRUDE PROTEIN, min	20.0 %
LYSINE, min	1.4 %
CRUDE FAT, min	7.0 %
CRUDE FIBER, max	4.5 %
CALCIUM, min	0.7 %
CALCIUM, max	1.2 %
PHOSPHORUS, min	0.6 %
SALT, min	0.4 %
SALT, max	0.9 %
SELENIUM, min	0.3 ppm
ZINC, min	4,150 ppm
PHYTASE (<i>Aspergillus niger</i>), min	209 FYT/lb*

*One unit of phytase activity (FYT) is defined as the quantity of enzyme which liberates 1 micromol of inorganic phosphorus per minute. Environmental factors such as feed pH, moisture, and processing methods such as pelleting or extrusion, may affect enzyme activity.

INGREDIENTS

Grain products, plant protein products, animal protein products, pork fat (preserved with BHA), roughage products, vegetable oil, dried fermentation biomass, calcium carbonate, sodium sulfite, dicalcium phosphate, monocalcium phosphate, L-lysine, salt, inulin, sodium bentonite, zinc oxide, methionine hydroxy analog, L-threonine, dried *Bacillus subtilis* fermentation product, chromium propionate, dried yeast fermentation solubles, sodium selenite, manganous oxide, dried *Bacillus licheniformis* fermentation product, monoglycerides, diglycerides, natural and artificial flavor, processed grain by-products, d-calcium pantothenate, calcium silicate, BHA (a preservative), disodium EDTA, silicon dioxide, lactic acid, choline chloride, phosphoric acid, menadione sodium bisulfite complex (source of vitamin K activity), vitamin A supplement, vitamin D supplement, niacin supplement, folic acid, biotin, yucca schidigera extract, copper sulfate, ferrous sulfate, tryptophan, vitamin E supplement, ethoxyquin, vitamin B12 supplement, riboflavin supplement, citric acid, thiamine mononitrate, dried *Aspergillus niger* fermentation products (phytase), hydrated sodium calcium aluminosilicate, propionic acid (a preservative), amorphous silicon dioxide, ammonium hydroxide, magnesium mica, hydrolyzed yeast, yeast extract, ethylenediamine dihydroiodide, selenium yeast

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

This product has been formulated specifically for swine and is not intended for other species.

FEEDING DIRECTIONS: Feed continuously as the sole ration for 14 days.

Feed as the sole ration to pigs from 26 to 40 days of age at the rate of 0.5 pound per 10 pounds body weight to

provide 10 milligrams per pound of Chlortetracycline per pound of body weight per day. A constant supply of fresh, clean drinking water is also necessary.

WARNING: Withdraw medicated feed 2 days before slaughter.

CAUTION: Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

SPECIAL CONSIDERATION: Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104oF (40oC), for more than 7 days.
LIMITATIONS FOR USE: Feed continuously as sole ration for 14 days. Feed approximately 400 g/ton Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb of bodyweight per day.

Net Weight 50 lb (22.68 kg) or bulk invoice.